CORRECTION Open Access

## Correction: Modified posterior approach of the knee in patients with diffuse pigmented Villonodular synovitis: case series of a single Institution's experience

Yi-Ping Wei and Shan-Wei Yang\*

Correction: BMC Musculoskelet Disord 23, 197 (2022) https://doi.org/10.1186/s12891-022-05103-4

Following the publication of the original article [1] the authors reported that a certain part of Table 2 has been mistyped.

The original article [1] has been updated. See page 2 for the corrected part of the table.

The original article can be found online at https://doi.org/10.1186/s12891-022-05103-4.

\*Correspondence: yangshanwei@yahoo.com.tw

Department of Orthopaedic, Kaohsiung Veterans General Hospital, 386, Ta-Chung 1st Rd, Kaohsiung, Taiwan, Republic of China



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/loublicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data

Table 2 Group comparison

Characteristics	SP group $(n = 13)$	DP group $(n=7)$	p value
Age (years)	39.62±15.78	45.86 ± 14.49	0.211 a
Gender (male/female)	6/7	6/1	0.158 b
Combined anterior incision (n)	12 (92.3%)	7 (100%)	1.000 b
Combined adjuvant radiotherapy (n)	8 (61.5%)	6 (85.7%)	0.354 <sup>b</sup>
Operation time (minutes)	115.00 ± 43.83 (PL only: 40; Anterior + PL: 100; Anterior + PM: 180; Anterior + PM + PL: 117.50 ± 37.88)	184.57 ± 68.59	0.019 <sup>a,c</sup>
Blood loss (ml)	$18.46 \pm 11.97$ (PL only: 5; Anterior + PL: 10; Anterior + PM: 20; Anterior + PM + PL: $20.50 \pm 12.57$ )	47.14±33.52	0.109 <sup>a,c</sup>
Postoperative WOMAC	91.23 ± 7.20 (75-99)	$76.00 \pm 16.57$ (48-98)	0.037 <sup>a</sup>
Postoperative MSTS	$24.23 \pm 2.68 (20-28)$	$22.43 \pm 4.69 (14-27)$	0.485 a
Recurrence (n)	2 (15.4%)	2 (28.6%)	0.587 <sup>b</sup>
Preoperative active ROM of the knee (degree)	$104.23 \pm 16.56$	$93.57 \pm 28.09$	0.438 <sup>a</sup>
Preoperative extension (degree)	$1.54 \pm 3.76$	$0\pm0$	0.588 a
Preoperative flexion (degree)	$105.77 \pm 14.41$	$93.57 \pm 28.09$	0.351 <sup>a</sup>
Postoperative active ROM of the knee (degree)	$122.31 \pm 14.52$	$100.71 \pm 23.53$	0.046 a
Postoperative extension (degree)	$2.31 \pm 4.39$ 124.62 ±12.98	$4.29 \pm 5.35$	0.485 a
Postoperative flexion (degree)	124.02 ±12.98 121.15 ± 15.70	$105.00 \pm 19.37$	-0.081 a
Postoperative ROM > 120° flexion (n)	<del>8 (61.5%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%) 9 (69.2%)</del>	3 (42.9%)	-0.642 b

Continuous data (i.e., age, operation time, blood loss, WOMAC, MSTS, and ROM) are presented as mean  $\pm$  SD (range); categorical data are presented as number (percentage)

Published online: 06 June 2022

## Reference

 Wei YP, Yang SW. Modified posterior approach of the knee in patients with diffuse pigmented Villonodular synovitis: case series of a single Institution's experience. BMC Musculoskelet Disord. 2022;23:197. https://doi.org/10.1186/s12891-022-05103-4.

## Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- $\bullet\,$  thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

## At BMC, research is always in progress.

**Learn more** biomedcentral.com/submissions



<sup>&</sup>lt;sup>a</sup> Mann–Whitney U test for comparing the continuous data from the two groups

<sup>&</sup>lt;sup>b</sup> Fisher's exact test for categorical data

<sup>&</sup>lt;sup>c</sup>The operation time and blood loss was compared between 10 patients (who received anterior, PM and PL incisions in SP group) and 7 patients (who received anterior and DP incisions in DP group)