

RESEARCH

Open Access



A multicenter study of 1-year mortality and walking capacity after spinal fusion surgery for cervical fracture in elderly patients

Takeshi Sasagawa^{1,2}, Noriaki Yokogawa¹, Hiroyuki Hayashi^{1,3}, Hiroyuki Tsuchiya¹, Kei Ando⁴, Hiroaki Nakashima⁴, Naoki Segi⁴, Kota Watanabe⁵, Satoshi Nori⁵, Kazuki Takeda^{5,6}, Takeo Furuya⁷, Atsushi Yunde⁷, Shota Ikegami⁸, Masashi Uehara⁸, Hidenori Suzuki⁹, Yasuaki Imajo⁹, Toru Funayama¹⁰, Fumihiko Eto¹¹, Akihiro Yamaji¹², Ko Hashimoto¹³, Yoshito Onoda¹³, Kenichiro Kakutani¹⁴, Yuji Kakiuchi¹⁴, Nobuyuki Suzuki¹⁵, Kenji Kato¹⁵, Yoshinori Terashima^{16,17}, Ryosuke Hirota¹⁶, Tomohiro Yamada^{18,19}, Tomohiko Hasegawa¹⁸, Kenichi Kawaguchi²⁰, Yohei Haruta²⁰, Shoji Seki²¹, Hitoshi Tonomura²², Munehiro Sakata^{22,23}, Hiroshi Uei^{24,25}, Hirokatsu Sawada²⁵, Hiroyuki Tominaga²⁶, Hiroto Tokumoto²⁶, Takashi Kaito²⁷, Yoichi Iizuka²⁸, Eiji Takasawa²⁸, Yasushi Oshima²⁹, Hidetomi Terai³⁰, Koji Tamai³⁰, Bungo Otsuki³¹, Masashi Miyazaki³², Hideaki Nakajima³³, Kazuo Nakanishi³⁴, Kosuke Misaki³⁴, Gen Inoue³⁵, Katsuhito Kiyasu³⁶, Koji Akeda³⁷, Norihiko Takegami³⁷, Toshitaka Yoshii³⁸, Masayuki Ishihara³⁹, Seiji Okada²⁷, Yasuchika Aoki⁴⁰, Katsumi Harimaya⁴¹, Hideki Murakami¹⁵, Ken Ishii^{42,43,44}, Seiji Ohtori⁷, Shiro Imagama⁴ and Satoshi Kato^{1*}

Abstract

Background: The 1-year mortality and functional prognoses of patients who received surgery for cervical trauma in the elderly remains unclear. The aim of this study is to investigate the rates of, and factors associated with mortality and the deterioration in walking capacity occurring 1 year after spinal fusion surgery for cervical fractures in patients 65 years of age or older.

Methods: Three hundred thirteen patients aged 65 years or more with a traumatic cervical fracture who received spinal fusion surgery were enrolled. The patients were divided into a survival group and a mortality group, or a maintained walking capacity group and a deteriorated walking capacity group. We compared patients' backgrounds, trauma, and surgical parameters between the two groups. To identify factors associated with mortality or a deteriorated walking capacity 1 year postoperatively, a multivariate logistic regression analysis was conducted.

Results: One year postoperatively, the rate of mortality was 8%. A higher Charlson comorbidity index (CCI) score, a more severe the American Spinal Cord Injury Association impairment scale (AIS), and longer surgical time were identified as independent factors associated with an increase in 1-year mortality. The rate of deterioration in walking capacity between pre-trauma and 1 year postoperatively was 33%. A more severe AIS, lower albumin (Alb) and hemoglobin (Hb) values, and a larger number of fused segments were identified as independent factors associated with the increased risk of deteriorated walking capacity 1 year postoperatively.

*Correspondence: skato323@gmail.com

¹ Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kanazawa University, 13-1 Takara-machi, Kanazawa, Ishikawa 920-8641, Japan
Full list of author information is available at the end of the article



Conclusions: The 1-year rate of mortality after spinal fusion surgery for cervical fracture in patients 65 years of age or older was 8%, and its associated factors were a higher CCI score, a more severe AIS, and a longer surgical time. The rate of deterioration in walking capacity was 33%, and its associated factors were a more severe AIS, lower Alb, lower Hb values, and a larger number of fused segments.

Keywords: Cervical fracture, Spinal fusion surgery, Elderly, Mortality, Walking capacity

Background

In step with the overall aging of the population, the frequency of occurrence of cervical spinal fractures in the elderly has increased in recent years [1, 2]. Cervical spine fractures may occur in isolation or in conjunction with a spinal cord injury and are relatively common among older adults, whose susceptibility may increase with minor trauma. Cervical spine fractures represent an important cause of mortality among adults aged 65 years or older [1, 2]. The mortality associated with cervical spinal fractures in elderly patients exceeds that in younger patients [3]. Compared with a hip fracture, a common type of fracture in the elderly, patients with cervical fractures had a greater mortality than those with hip fractures [1]. Surgical treatment for cervical trauma is a more invasive option compared with conservative treatment. Therefore, it is important to understand mortality and functional outcomes after surgical treatment for cervical trauma.

In previous reports that have explored the relationship between mortality and cervical spinal fractures in the elderly, only in-hospital mortality has been discussed [2–5], and only a few reports have referred to 1 year mortality [6, 7]. Furthermore, most of the previous reports included both surgical and conservative treatments [2–6], and few reports specify the mortality of patients treated with surgery alone [7]. The 1-year mortality of patients who received surgery for cervical trauma remains unclear. Moreover, to our knowledge, although functional outcomes after surgery for cervical trauma are important, 1-year postoperative functional prognoses remain unreported. The aim of this study is to investigate the rates of, and factors associated with mortality and any changes in walking capacity occurring 1 year after spinal fusion surgery for cervical fractures in patients 65 years of age or older.

Materials and methods

Study design and ethical considerations

This study retrospectively analyzed multicenter registry data collected by the Japan Association of Spine Surgeons with Ambition (JASA). The institutional review board of the representative facility reviewed and approved this study. No funds were received in support of this study. No benefits in any form have been or will be received

from a commercial party related directly or indirectly to the subject of this manuscript.

Patient population

Patients aged 65 years or more with a traumatic cervical fracture who received spinal fusion surgery from February 2010 to August 2019 at 68 institutions registered with JASA were considered for inclusion in this study. Patients who received surgery more than 90 days postinjury, or those for whom missing data pertaining to the type of surgery and level(s) of fusion were missing, were excluded. Of the 418 patients who were eligible for study participation, 105 of them were lost to follow-up 1 year postoperatively (follow-up rate was 75%) and 313 patients were enrolled.

Data collection

Patients' background data

Data for each patient, including age at time of injury, gender, height, weight, pre-trauma walking capacity, Charlson Comorbidity Index (CCI) score [8], and blood test results at the first preoperative visit were collected. We defined the main walking style in everyday life as walking capacity. Patients' walking capacities were divided into four grades: independent, able to walk with a T-cane, able to walk with a walker, or inability to walk. Blood tests were used to measure albumin (Alb) and hemoglobin (Hb).

Trauma data

Collected radiographic data included the fracture level(s), presence of facet interlocking, and comorbid major organ injury. Comorbid major organ injury was defined as other trauma requiring surgery, hemothorax requiring a chest drain, or brain injury with consciousness disturbance. Neurological impairment was assessed using the American Spinal Cord Injury Association (ASIA) impairment scale (AIS), ranging from Grade A (complete impairment) to Grade E (normal function) [9]. In patients with AIS A to D, spinal cord injury type (transverse or central) was also investigated.

Surgical data

The surgical data documented included the surgical approach (anterior, posterior, or combined), number of

fused levels, presence of occipitocervical fusion, surgical bleeding (mL), and surgical time (min).

Operative outcomes

Collected operative outcomes included intraoperative and postoperative complications, patient mortality, and walking capacity 1 year postoperatively.

Statistical analysis

All 313 patients were divided into a survival group and a mortality group. We also divided the patients who were able to walk before injury and survived 1 year postoperatively into a 'maintained walking capacity' group and a 'deteriorated walking capacity' group. Deteriorated walking capacity applied when a patient's walking capacity decreased by at least one grade between pre-trauma and 1 year postoperatively. We compared patients' backgrounds, trauma, and surgical parameters between the two groups. All data are expressed as mean \pm standard deviation. A Mann–Whitney U test, chi square test, or Fisher's exact test was used to compare each item. To identify factors associated with mortality or a

deteriorated walking capacity 1 year postoperatively, a multivariate logistic regression analysis was conducted in which mortality or a deteriorated walking capacity were used as a dependent variable. Items that were significantly different by univariate analysis were independent variables. Differences were considered significant at $P < 0.05$. All statistical analyses were performed using IBM SPSS Statistics for Windows (version 22; IBM Corp, Armonk, NY).

Results

The demographic data for 313 cases are presented in Table 1. There were 201 men and 112 women of mean age 74.6 years in this study. Of the 313 patients we assessed, 99% were able to walk before injury. The most common site for cervical fractures was C6–7 (46%), followed by C3–5 (43%), and C1–2 (31%). The frequency of facet interlocking was 28%, and a spinal cord injury occurred in 51% of patients. The type of spinal cord injury in 158 patients with AIS A–D was transverse in 88 patients (56%) and central in 70 patients (44%). Surgery was most frequently via the posterior approach (87%).

Table 1 Patient characteristics and demographics

Parameter ^a	n	N (%) ^b
Patients, n		313
Age, years		74.6 \pm 6.2
Gender (male: female), n		201:112
Height, cm	298	158.9 \pm 9.7
Weight, kg	300	55.8 \pm 10.3
Pre-trauma walking capacity, n		
Independent / T-cane / walker / inability to walk		291 / 13 / 5 / 4
Charlson comorbidity index	307	0.6 \pm 1.0
Blood test data, g/dL		
Albumin	267	3.7 \pm 0.5
Hemoglobin	309	12.6 \pm 1.9
Level of fracture, n (%)		
C1–2 / C3–5 / C6–7		98 (31) / 113 (36) / 144 (46)
Facet interlocking, n (%)		89 (28)
ASIA Impairment Scale, n (%)		
A / B / C / D / E	312	27 (9) / 14 (5) / 43 (14) / 74 (24) / 154 (49)
Comorbid major organ injury, n (%)		31 (10)
Surgery		
Anterior / Posterior / Combined, n (%)		30 (10) / 273 (87) / 10 (3)
Number of fused segments		2.4 \pm 1.8
Occipitocervical fusion, n (%)		16 (5)
Surgical bleeding, mL	291	174 \pm 71
Surgical time, min	288	247 \pm 387

ASIA American Spinal Injury Association

^a Data are the mean \pm standard deviation unless otherwise shown

^b Results for a denominator of $N = 313$, unless otherwise indicated by n in middle column

The intraoperative and postoperative complications of the 313 cases are presented in Table 2. The most frequent intraoperative complication was an iatrogenic dural tear (2%), and the most frequent postoperative complication was pneumonia (11%).

Life prognosis

One year postoperatively, 25 out of 313 patients had died and the rate of mortality was 8% (Table 2). Death occurred within one month in 12 of 25 patients (48%) and within six months in 20 of 25 patients (80%). The causes of death were respiratory complications (respiratory failure, asphyxia, pneumonia) in 12 cases, cerebral infarction in 3 cases, gastrointestinal complications in 2 cases, malignancy in 2 cases, heart failure in 2 cases, pulmonary embolism in 1 case, massive intraoperative bleeding in 1 case, and details unknown in 2 cases. A comparison between the survival group ($n=288$) and the mortality group ($n=25$) revealed that being male, being assessed as having a higher CCI score, a severe AIS, and/or a longer surgical time were more significantly associated with the mortality group than the survival group (Table 3). A multivariate logistic regression analysis was conducted to identify factors associated with 1-year mortality. A higher CCI score (odds ratio [OR] = 2.046, 95% confidence interval [CI]: 1.398–2.993), more severe AIS

(OR = 2.205, 95% CI: 1.586–3.065), and longer surgical time (OR = 1.009, 95% CI: 1.002–1.015) were identified as independent factors associated with an increase in 1-year mortality (Table 4).

Walking capacity

Of 313 patients, 284 patients were able to walk before injury and survived 1 year postoperatively. Among these 284 patients, 93 patients (33%) exhibited a deterioration in their pre-trauma walking capacity 1 year postoperatively (Table 2). The rate of walking capacity deterioration 1-year postoperatively by AIS was 100% for AIS A/B (all cases with transverse spinal cord injuries), 63% (transverse type 72%, central type 47%) for AIS C, 21% (transverse type 44%, central type 13%) for AIS D, and 18% for AIS E. A comparison between the maintained walking capacity group ($n=191$) and the deteriorated walking capacity group ($n=93$) revealed the values of Alb and Hb, and the frequency of C1–2 fractures were significantly less in the deteriorated walking capacity group. In contrast, the frequency of C3–5 fractures, severe AIS, number of fused segments, and surgical bleeding were significantly higher in the deteriorated walking capacity group than the maintained walking capacity group (Table 5). A multivariate logistic regression analysis was conducted to identify factors associated with deteriorated walking capacity 1 year postoperatively. A more severe AIS (OR = 4.092, 95% CI: 2.767–6.052), lower Alb (OR = 0.341, 95% CI: 0.172–0.672), lower Hb (OR = 0.797, 95% CI: 0.660–0.961) values and a larger number of fused segments (OR = 1.241, 95% CI: 1.023–1.506) were identified as independent factors associated with the increased risk of deteriorated walking capacity 1 year postoperatively (Table 6).

Table 2 Overall operative outcomes

Outcomes	N = 313
Intraoperative complications, n (%)	
Dural tear	7 (2)
Vertebral artery injury	4 (1)
Spinal cord injury	1 (< 1)
Postoperative complications, n (%)	
Pneumonia	33 (11)
Delirium	30 (10)
Urinary tract infection	27 (9)
Surgical site infection	6 (2)
Cerebral infarction	5 (2)
Instrumentation failure	4 (1)
Cerebrospinal fluid leakage	4 (1)
Pulmonary embolism	3 (1)
Epidural hematoma	1 (< 1)
1-year mortality, %	
Overall	8
ASIA Impairment Scale A / B / C / D / E	44 / 14 / 5 / 1 / 5
Walking capacity deterioration 1-year postoperatively ^a , %	
Overall	33
ASIA Impairment Scale A / B / C / D / E	100 / 100 / 63 / 21 / 18

ASIA American Spinal Injury Association

^a $n=284$ patients who can walk before injury

Discussion

Based on the present study, we report a 1-year mortality rate of 8% after spinal fusion surgery for cervical fractures in patients 65 years of age or older. In previous reports of cervical spinal fractures in the elderly, the rates of mortality coexistent with a spinal cord injury were 7–53% [2–6]. In our study, 51% of cervical fractures were associated with a spinal cord injury. Thus, our results reflect cervical fractures with a high rate of concomitant spinal cord injury. This can be attributed to the fact that the present study comprised patients who received spinal fusion surgery. Previous reports of mortality associated with cervical spine fractures in the elderly have referred to an in-hospital mortality rate of 8–14% [2–5], and a 1-year mortality rate of 28–29% [6, 7]. Most of those previous reports included both surgical and conservative treatments [2–6]. Although Sander et al. have reported on postsurgical mortality, they noted decompression

Table 3 Comparison of admission data between survival and mortality groups 1 year postoperatively

	n	Survival group (n = 288)	n	Mortality group (n = 25)	P
Patient background					
Age, years		74.5 ± 6.3		76.6 ± 5.4	0.07
Gender (male / female), n		179 / 109		22 / 3	0.01*
Height, cm	274	158.7 ± 9.7	24	161.5 ± 9.9	0.19
Weight, kg	277	55.7 ± 10.3	24	56.8 ± 10.8	0.78
Pre-trauma walking capacity, n					
Independent / T-cane / walker / inability to walk		269 / 12 / 3 / 4		22 / 1 / 2 / 0	0.14
Charlson comorbidity index	282	0.5 ± 1.0		1.3 ± 1.6	< 0.01*
Blood test data, g / dL					
Albumin	247	3.7 ± 0.5	20	3.6 ± 0.4	0.16
Hemoglobin	285	12.7 ± 1.9	24	12.1 ± 2.3	0.33
Level of fracture, n (%)					
C1–2		93 (32)		5 (20)	0.20
C3–5		102 (35)		11 (44)	0.39
C6–7		130 (45)		14 (56)	0.30
Facet interlocking, n (%)		81 (28)		8 (32)	0.68
ASIA Impairment Scale, n (%)					
A / B / C / D / E	287	15 (5) / 12 (4) / 41 (14) / 73 (25) / 146 (51)		12 (48) / 2 (8) / 2 (8) / 1 (4) / 8 (32)	< 0.01*
Comorbid major organ injury, n (%)					
		30 (10)		1 (4)	0.26
Surgery					
Anterior / Posterior / Combined, n (%)		28 (10) / 251 (87) / 9 (3)		2 (8) / 22 (88) / 1 (4)	0.94
Number of fused segments		2.3 ± 1.7		3.0 ± 2.0	0.10
Occipitocervical fusion, n (%)		15 (5)		1 (4)	0.63
Surgical bleeding, mL	266	228 ± 304	22	481 ± 886	0.10
Surgical time, min	268	171 ± 68	23	217 ± 83	< 0.01*

ASIA American Spinal Injury Association

^a Data are the mean ± standard deviation unless otherwise shown

^b Results for denominator n = 288 (survivor group) or n = 25 (mortality group), unless otherwise indicated in the column to the left of each of these group results, respectively

* P < 0.05

Table 4 Multivariate logistic regression analysis of associated factors of mortality 1 year postoperatively

Variables	OR	95% CI	P
Gender: male	0.419	0.109–1.611	0.21
Charlson comorbidity index	2.046	1.398–2.993	< 0.01*
ASIA Impairment Scale	2.205	1.586–3.065	< 0.01*
Surgical time (min)	1.009	1.002–1.015	0.01*

OR Odds ratio, CI Confidence interval, ASIA American Spinal Injury Association

The ASIA Impairment Scale is treated as an ordinal variable (A-E)

* P < 0.05

surgery was included in addition to fusion surgery [7]. In contrast, our study comprised only patients who received spinal fusion surgery for cervical fractures. The best studied relationship between injury type and mortality rate is for hip fractures. The reported 1-year mortality rate after

hip fracture in the elderly is 10–30% [10, 11]. Patients with a cervical fracture incur a greater risk of mortality compared with those who sustain a hip fracture [1]. However, we did not demonstrate this phenomenon in the current study.

Our study did show that a higher CCI score, more severe AIS, and a longer surgical time were identified as independent factors associated with increasing 1-year mortality. Previously, whether a greater number of comorbidities was associated with mortality or not is controversial in the elderly with cervical fractures [6, 12]. The results of this study indicated that a greater number of comorbidities was associated with an increased mortality risk in patients treated surgically. Previously, a neurological deficit has been linked to mortality after a cervical fracture in the elderly [1–3, 6, 13] and mortality has been correlated with the severity of a neurological deficit [13, 14]. The results of our

Table 5 Comparison of admission data between maintained and deteriorated walking capacity groups 1 year postoperatively

	n	Maintained walking capacity group (n = 191)	n	Deteriorated walking capacity group (n = 93)	P
Patient background					
Age, years		74.3 ± 6.2		74.6 ± 6.4	0.78
Gender (male / female), n		111 / 80		64 / 29	0.08
Height, cm	185	158.4 ± 9.7	85	159.3 ± 9.6	0.36
Weight, kg	186	55.0 ± 10.0	87	57.5 ± 10.9	0.07
Pre-trauma walking capacity, n					
Independent / T-cane / walker		183 / 6 / 2		86 / 6 / 1	0.35
Charlson comorbidity index	189	0.5 ± 1.0	89	0.6 ± 0.9	0.20
Blood test data, g / dL					
Albumin	156	3.8 ± 0.5	87	3.6 ± 0.6	< 0.01*
Hemoglobin	188	12.9 ± 1.9	84	12.3 ± 1.8	0.02*
Level of fracture, n (%)					
C1–2		71 (37)		22 (24)	0.02*
C3–5		59 (31)		41 (44)	0.03*
C6–7		86 (45)		42 (45)	0.98
Facet interlocking, n (%)		53 (28)		28 (30)	0.68
ASIA Impairment Scale, n (%)					
A / B / C / D / E		0 / 0 / 15 (8) / 57 (30) / 119 (62)		15 (16) / 11 (12) / 25 (27) / 15 (16) / 27 (29)	< 0.01*
Comorbid major organ injury, n (%)		16 (8)		14 (15)	0.09
Surgery					
Anterior / Posterior / Combined, n (%)		20 (10) / 164 (86) / 7 (4)		7 (8) / 84 (90) / 2 (2)	0.56
Number of fused segments		2.1 ± 1.6		2.6 ± 1.9	0.01*
Occipitocervical fusion, n (%)		8 (4)		7 (8)	0.18
Surgical bleeding, mL	176	187 ± 220	86	291 ± 363	0.01*
Surgical time, min	181	168 ± 72	83	176 ± 60	0.25
Walking capacity 1 year postoperatively, n					
Independent / T-cane / walker / inability to walk		184 / 6 / 1 / 0		0 / 32 / 33 / 28	

^a Data are the mean ± standard deviation unless otherwise shown

^b Results for denominator of n = 191 (maintained walking capacity group) or n = 93 (deteriorated walking capacity group), unless otherwise indicated by in the column to the left of each of these group results, respectively

ASIA American Spinal Injury Association

* P < 0.05

Table 6 Multivariate logistic regression analysis of associated factors of deteriorated walking capacity 1 year postoperatively

Variables	OR	95% CI	P
Albumin (g/dL)	0.341	0.172 – 0.672	< 0.01*
Hemoglobin (g/dL)	0.797	0.660 – 0.961	0.02*
Fracture at C1–2	0.489	0.210 – 1.140	0.10
Fracture at C3–5	0.263	0.323 – 1.362	0.26
ASIA Impairment Scale	4.092	2.767 – 6.052	< 0.01*
Fused segments	1.241	1.023 – 1.506	0.03
Surgical bleeding (mL)	1.001	1.000 – 1.002	0.19

OR Odds ratio, CI Confidence interval, ASIA American Spinal Injury Association
The ASIA Impairment Scale is treated as an ordinal variable (A-E)

* P < 0.05

study supported the same finding. Daneshvar et al. demonstrated that an injury at or above C4 had a 7.1 times higher risk of mortality compared with injuries below C4 when spinal cord injuries were related to cervical spine fractures [13]. The logical connection to consider is that the more severe the spinal cord injury, the greater the impact on respiratory muscle function and hence an increased risk for mortality.

We included surgical factors in our investigation because our study comprised patients who were treated surgically. As a result, a longer surgical time was identified as an independent factor associated with increasing 1-year mortality. There are two possible explanations. First, a cervical fracture requiring a long surgical time

is typically indicative of severe trauma leading to poor general condition. Second, based on our result that 48% of those in the mortality group died within one month postoperatively, surgical invasiveness is considered to be related to mortality in the immediate postoperative window. Some reports indicate that surgical invasiveness for spinal trauma in the acute phase is related to the mortality [6, 15]. Also, greater surgical invasiveness during spinal surgery increases postoperative complications, particularly in the elderly [16]. In our study results pneumonia was the most frequent postoperative complication reported. In addition, respiratory complications were the most common cause of death. Bokhari et al. reported that the occurrence of pneumonia is more frequent during surgical treatment than conservative treatment in elderly patients with cervical fractures [17]. In short, there is a possibility that surgical invasiveness gave rise to a decline in the general condition of the patients, resulting in poor respiratory function leading to mortality. In terms of surgical factors, in our study we were also able to demonstrate that surgical time was a more important factor than surgical approach, surgical bleeding, or number of fused levels.

The rate of deterioration in walking capacity between the time of injury and 1 year postoperatively was 33% in this study. To our knowledge, there are no reports of functional outcomes 1 year after spinal fusion surgery for elderly patients with cervical fractures. With regard to hip fractures in the elderly, the reported 1-year postoperative rate of deterioration in walking capacity is 26–61% [18–20]. Therefore, the rate of deterioration in walking capacity in the elderly with a cervical fracture in our study was comparable to that reported for hip fracture.

A more severe AIS, lower Alb, lower Hb values and a larger number of fused segments were identified as independent factors associated with an increased risk of deteriorated walking capacity 1 year postoperatively. Previously, neurological deficits have been related to poor functional outcomes [5]. Our study also demonstrated that the severity of neurological deficits was related to poor functional outcomes for as long as 1 year postoperatively. Reports have indicated that recovery from a spinal cord injury is poor in the elderly population [13]. Moreover, even when neurological function recovered, elderly patients experience difficulties translating improvements in a neurological outcome into functional changes [21]. For these reasons, the severity of a neurological deficit at the time of injury is considered to be strongly associated with a deterioration in walking capacity 1 year postoperatively.

Serum Alb has traditionally been used as a marker for poor health and nutrition. In the elderly with hip fractures, hypoalbuminemia or poor nutrition is related to

poor functional outcomes [22, 23]. Hypoalbuminemia also has been reported to be associated with poor functional outcomes in cervical spinal cord injury [24]. In short hypoalbuminemia is considered to lead to poor recovery of neuromuscular function. We demonstrated that low preoperative Alb values were associated with a poor functional outcome in the elderly with a cervical fracture, as well.

A low level of preoperative Hb has been associated with a poor functional outcome according to some reports of hip fractures in the elderly [25, 26]. Low preoperative Hb values were associated with poor functional outcomes in the elderly with cervical fracture in our study also. A possible explanation for the relationship between low Hb values and poor functional outcomes is that Hb status could be a marker of an underlying comorbidity [25]. Anemia has been observed to be a risk factor for the frailty phenotype in the elderly [27]. Thus, low Hb values preoperatively could be a reflection of greater frailty associated with a poor functional outcome [25].

A larger number of fused segments was identified as an independent factor associated with the increased risk of a deteriorated walking capacity one year postoperatively. Previously, it was reported that a larger number of fused segments was associated with disability and compromises to activities of daily living [28]. Furthermore, whole cervical spine fixation increases stride time and decreases stride length. Fixation reduces motions between the shoulder girdle and the trunk, and between the trunk and the pelvis, and decreases hip motion [29]. In short, the increasing limitation of range of motion of the cervical spine impacts walking capacity. Although a limitation of this study is that the range of motion of the cervical spine was not assessed, it is possible in elderly patients with cervical fractures that limitation of the range of cervical spine motion due to fusion of more segments impacts walking capacity.

There are some possible limitations in the present study. Because of its retrospective design, there are some missing data. Because this was a retrospective multicenter study, the indications for surgery, choice of surgical technique, and postoperative treatment were left to the discretion of the surgeon at each hospital. Lack of detailed information such as surgeon, screw type, anesthesia and geriatric medical care, is also a limitation of this study. Although the follow-up rate (75%) was high, a selection bias could occur due to the retrospective investigation of only those patients who could be followed-up for 1 year. Because the sample size of the mortality group was small, a further study using a larger sample size is needed to better understand the factors that were associated with mortality after spinal fusion surgery for a cervical fracture in the elderly.

Conclusions

The 1-year rate of mortality after spinal fusion surgery for cervical fracture in patients 65 years of age or older was 8%. A higher CCI score, a more severe AIS, and a longer surgical time were identified as independent factors associated with increasing 1-year mortality. The rate of deterioration in walking capacity between pre-trauma and 1 year postoperatively was 33%. A more severe AIS, lower Alb, lower Hb values, and a larger number of fused segments were identified as independent factors associated with an increasing risk of deterioration in walking capacity 1 year postoperatively.

Abbreviations

JASA: Japan Association of Spine Surgeons with Ambition; CCI: Charlson comorbidity index; ASIA: American Spinal Cord Injury Association; AIS: American Spinal Cord Injury Association impairment scale; Alb: Albumin; Hb: Hemoglobin; OR: Odds ratio; CI: Confidence interval.

Acknowledgements

Not applicable.

Authors' contributions

Research conception and design: SK, TS. Data collection: TS, NY, HH, H Tsuchiya, K Ando, H Nakashima, N Segi, KW, SN, K Takeda, T Furuya, A Yunde, S Ikegami, MU, H Suzuki, Y Imajo, T Funayama, FE, A Yamaji, K Hashimoto, Y Onoda, K Kakutani, YK, N Suzuki, K Kato, YT, RH, T Yamada, TH, K Kawaguchi, YH, SS, H Tonomura, MS, HU, H Sawada, H Tominaga, H Tokumoto, TK, Y Iizuka, ET, Y Oshima, H Terai, K Tamai, BO, MM, H Nakajima, KN, KM, GI, K Kiyasu, K Aakeda, NT, T Yoshii, MI, S Okada, YA, K Harimaya, HM, KI, S Ohtori, S Imagama, SK. Interpretation of data: SK, TS, NY. Statistical analysis: SK, TS, NY. Drafting the manuscript: TS. Manuscript review: SK, NY. Study supervision: K Ando, KW, T Furuya, TK, GI. Approval of the final manuscript: all the authors listed above.

Funding

No funding.

Availability of data and materials

The datasets during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The study has been performed in accordance with the ethical standards in the 1964 Declaration of Helsinki. The institutional review board of the representative facility (Kanazawa University) reviewed and approved this study (No. 3352-1).

All subjects gave written informed consent to participate in the study.

Consent for publication

Not applicable.

Competing interests

The authors have no conflicts of interest to declare.

Author details

¹Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kanazawa University, 13-1 Takara-machi, Kanazawa, Ishikawa 920-8641, Japan.

²Department of Orthopaedic Surgery, Toyama Prefectural Central Hospital, 2-2-78 Nishinagae, Toyama, Toyama 930-8550, Japan.

³Department of Orthopaedic Surgery, Tonami General Hospital, Shintomicho 1-61, Tonami, Toyama 939-1395, Japan.

⁴Department of Orthopaedic Surgery, Graduate School of Medicine, Nagoya University, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan.

⁵Department of Orthopaedic Surgery, Keio University School of Medicine,

35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan.

⁶Department of Orthopaedic Surgery, Japanese Red Cross Shizuoka Hospital, 8-2 Otemachi, Aoi-ku, Shizuoka 420-0853, Japan.

⁷Department of Orthopaedic Surgery, Graduate School of Medicine, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba, Chiba 260-8670, Japan.

⁸Department of Orthopaedic Surgery, Shinshu University School of Medicine, 3-1-1 Asahi, Matsumoto, Nagano 390-8621, Japan.

⁹Department of Orthopaedic Surgery, Yamaguchi University Graduate School of Medicine, 1-1-1 Minami-kogushi, Ube city, Yamaguchi 755-8505, Japan.

¹⁰Department of Orthopaedic Surgery, Faculty of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan.

¹¹Department of Orthopaedic Surgery, Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan.

¹²Department of Orthopaedic Surgery, Ibaraki Seinan Medical Center Hospital, 2190, Sakaimachi, Sashima, Ibaraki 306-0433, Japan.

¹³Department of Orthopaedic Surgery, Tohoku University Graduate School of Medicine, 1-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8574, Japan.

¹⁴Department of Orthopaedic Surgery, Kobe University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe 650-0017, Japan.

¹⁵Department of Orthopaedic Surgery, Nagoya City University Graduate School of Medical Sciences, 1 Kawasumi, Mizuho-cho, Mizuho-ku, Nagoya 467-8601, Japan.

¹⁶Department of Orthopaedic Surgery, Sapporo Medical University, South 1-West 16-291, Chuo-ku, Sapporo 060-8543, Japan.

¹⁷Department of Orthopaedic Surgery, Matsuda Orthopedic Memorial Hospital, North 18-East 4-1 Kita-ku, Sapporo 001-0018, Japan.

¹⁸Department of Orthopaedic Surgery, Hamamatsu University School of Medicine, 1-20-1, Handayama, Higashi-ku, Hamamatsu City, Shizuoka 431-3192, Japan.

¹⁹Department of Orthopaedic Surgery, Nagoya Kyoritsu Hospital, 1-172 Hokke, Nakagawa-ku, Nagoya-shi, Aichi 454-0933, Japan.

²⁰Department of Orthopaedic Surgery, Graduate School of Medical Sciences, Kyushu University, 3-1-1 Maidashi Higashi-ku, Fukuoka 812-8582, Japan.

²¹Department of Orthopaedic Surgery, Faculty of Medicine, University of Toyama, 2630 Sugitani, Toyama, Toyama 930-0194, Japan.

²²Department of Orthopaedics, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kawaramachi-Hirokoji, Kamigyo-ku, Kyoto 602-8566, Japan.

²³Department of Orthopaedics, Saiseikai Shiga Hospital, 2-4-1 Ohashi Ritto, Shiga 520-3046, Japan.

²⁴Department of Orthopaedic Surgery, Nihon University Hospital, 1-6 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-8393, Japan.

²⁵Department of Orthopaedic Surgery, Nihon University School of Medicine, 30-1 Oyaguchi Kami-cho, Itabashi-ku, Tokyo 173-8610, Japan.

²⁶Department of Orthopaedic Surgery, Graduate School of Medical and Dental Sciences, Kagoshima University, 8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan.

²⁷Department of Orthopaedic Surgery, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

²⁸Department of Orthopaedic Surgery, Graduate School of Medicine, Gunma University, 3-39-22 Showa, Maebashi, Gunma 371-8511, Japan.

²⁹Department of Orthopaedic Surgery, The University of Tokyo Hospital, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan.

³⁰Department of Orthopaedic Surgery, Osaka City University Graduate School of Medicine, 1-4-3 Asahimachi, Abeno-ku, Osaka-city, Osaka 545-8585, Japan.

³¹Department of Orthopaedic Surgery, Graduate School of Medicine, Kyoto University, 54 Shogoin-Kawaracho, Sakyo-ku, Kyoto, Japan.

³²Department of Orthopaedic Surgery, Faculty of Medicine, Oita University, 1-1 Idaigaoka, Hasama-machi, Yufu-shi, Oita 879-5593, Japan.

³³Department of Orthopaedics and Rehabilitation Medicine, Faculty of Medical Sciences, University of Fukui, 23-3 Matsuoka Shimoaizuki, Eiheiji-cho, Yoshida-gun, Fukui 910-1193, Japan.

³⁴Department of Orthopedics, Traumatology and Spine Surgery, Kawasaki Medical School, Matsushima, Kurashiki, Okayama 577701-0192, Japan.

³⁵Department of Orthopaedic Surgery, Kitasato University School of Medicine, 1-15-1, Kitazato, Minami-ku, Sagami-hara, Kanagawa 252-0374, Japan.

³⁶Department of Orthopaedic Surgery, Kochi Medical School, Kochi University, KohasuNankoku, Oko-cho 783-8505, Japan.

³⁷Department of Orthopaedic Surgery, Mie University Graduate School of Medicine, 2-174 Edobashi, Tsu city, Mie 514-8507, Japan.

³⁸Department of Orthopaedic Surgery, Tokyo Medical and Dental University, Yushima 1-5-45, Bunkyo-Ku, Tokyo 113-8519, Japan.

³⁹Department of Orthopaedic Surgery, Kansai Medical University Hospital, 2-3-1 Shinmachi, Hirakata, Osaka 573-1191, Japan.

⁴⁰Department of Orthopaedic Surgery, Eastern Chiba Medical Center, 3-6-2 Okayamadai, Togane, Chiba 283-8687, Japan.

⁴¹Department of Orthopaedic Surgery, Kyushu University Beppu Hospital, Tsurumibarui, Beppu, Oita 874-0838, Japan.

⁴²Department of Orthopaedic Surgery, School of Medicine, International University of Health and Welfare, 852 Hatakeda, Narita, Chiba 286-0124, Japan.

⁴³Department of Orthopaedic Surgery, International University of Health and Welfare Narita Hospital, 852 Hatakeda, Narita, Chiba 286-0124, Japan.

⁴⁴Department

of Orthopaedic Surgery and Spine and Spinal Cord Center, International University of Health and Welfare Mita Hospital, 1-4-3 Mita, Minato-ku, Tokyo 108-8329, Japan.

Received: 2 March 2022 Accepted: 9 August 2022

Published online: 20 August 2022

References

- Cooper Z, Mitchell SL, Lipsitz S, Harris MB, Ayanian JZ, Bernacki RE, et al. Mortality and readmission after cervical fracture from a fall in older adults: comparison with hip fracture using national medicare data. *J Am Geriatr Soc*. 2015;63(10):2036–42.
- Asemota AO, Ahmed AK, Purvis TE, Passias PG, Goodwin CR, Sciubba DM. Analysis of cervical spine injuries in elderly patients from 2001 to 2010 using a nationwide database: increasing incidence, overall mortality, and inpatient hospital charges. *World Neurosurg*. 2018;120:e114–30.
- Sokolowski MJ, Jackson AP, Haak MH, Meyer PR Jr, Sokolowski MS. Acute mortality and complications of cervical spine injuries in the elderly at a single tertiary care center. *J Spinal Disord Tech*. 2007;20(5):352–6.
- Menendez ME, Ring D, Harris MB, Cha TD. Predicting in-hospital mortality in elderly patients with cervical spine fractures: a comparison of the Charlson and Elixhauser comorbidity measures. *Spine (Phila Pa 1976)*. 2015;40(11):809–15.
- Malik SA, Murphy M, Connolly P, O'Byrne J. Evaluation of morbidity, mortality and outcome following cervical spine injuries in elderly patients. *Eur Spine J*. 2008;17(4):585–91.
- Harris MB, Reichmann WM, Bono CM, Bouchard K, Corbett KL, Warholic N, et al. Mortality in elderly patients after cervical spine fractures. *J Bone Joint Surg Am*. 2010;92(3):567–74.
- Sander AL, El Saman A, Delfosse P, Wutzler S, Meier S, Marzi I, et al. Cervical spine fractures in the elderly: morbidity and mortality after operative treatment. *Eur J Trauma Emerg Surg*. 2013;39(5):469–76.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–83.
- Kirshblum SC, Burns SP, Biering-Sorensen F, Donovan W, Graves DE, Jha A, et al. International standards for neurological classification of spinal cord injury (revised 2011). *J Spinal Cord Med*. 2011;34(6):535–46.
- Downey C, Kelly M, Quinlan JF. Changing trends in the mortality rate at 1-year post hip fracture - a systematic review. *World J Orthop*. 2019;10(3):166–75.
- Shigemoto K, Sawaguchi T, Horii T, Goshima K, Iwai S, Higashikawa T, et al. Multidisciplinary care model for geriatric patients with hip fracture in Japan: 5-year experience. *Arch Orthop Trauma*. 2022;142(9):2205–14.
- Golob JF Jr, Claridge JA, Yowler CJ, Como JJ, Peerless JR. Isolated cervical spine fractures in the elderly: a deadly injury. *J Trauma*. 2008;64(2):311–5.
- Daneshvar P, Roffey DM, Brikeet YA, Tsai EC, Bailey CS, Wai EK. Spinal cord injuries related to cervical spine fractures in elderly patients: factors affecting mortality. *Spine J*. 2013;13(8):862–6.
- Inglis T, Banaszek D, Rivers CS, Kurban D, Evaniew N, Fallah N, et al. In-Hospital Mortality for the Elderly with Acute Traumatic Spinal Cord Injury. *J Neurotrauma*. 2020;37(21):2332–42.
- Kerwin AJ, Frykberg ER, Schinco MA, Griffen MM, Arce CA, Nguyen TQ, et al. The effect of early surgical treatment of traumatic spine injuries on patient mortality. *J Trauma*. 2007;63(6):1308–13.
- Nagata K, Shinozaki T, Yamada K, Nakajima K, Nakamoto H, Yamakawa K, et al. A sliding scale to predict postoperative complications undergoing posterior spine surgery. *J Orthop Sci*. 2020;25(4):545–50.
- Bokhari AR, Sivakumar B, Sefton A, Lin JL, Smith MM, Gray R, et al. Morbidity and mortality in cervical spine injuries in the elderly. *ANZ J Surg*. 2019;89(4):412–7.
- Cooper C. The crippling consequences of fractures and their impact on quality of life. *Am J Med*. 1997;103(2A):12S–17S (discussion 17S–19S).
- Kitamura S, Hasegawa Y, Suzuki S, Sasaki R, Iwata H, Wingstrand H, et al. Functional outcome after hip fracture in Japan. *Clin Orthop Relat Res*. 1998;348:29–36.
- Moerman S, Mathijssen NM, Tuinebreijer WE, Nelissen RG, Vochteloo AJ. Less than one-third of hip fracture patients return to their prefracture level of instrumental activities of daily living in a prospective cohort study of 480 patients. *Geriatr Gerontol Int*. 2018;18(8):1244–8.
- Jakob W, Wirz M, van Hedel HJ, Dietz V, EM-SCI Study Group. Difficulty of elderly SCI subjects to translate motor recovery—"body function"—into daily living activities. *J Neurotrauma*. 2009;26(11):2037–44.
- Sim SD, Sim YE, Tay K, Howe TS, Png MA, Chang CCP, et al. Preoperative hypoalbuminemia: Poor functional outcomes and quality of life after hip fracture surgery. *Bone*. 2021;143: 115567.
- Lieberman D, Friger M, Lieberman D. Inpatient rehabilitation outcome after hip fracture surgery in elderly patients: a prospective cohort study of 946 patients. *Arch Phys Med Rehabil*. 2006;87(2):167–71.
- Jin GX, Li L, Cui SQ, Duan JZ, Wang H. Persistent hypoalbuminemia is a predictor of outcome in cervical spinal cord injury. *Spine J*. 2014;14(9):1902–8.
- Sim YE, Sim SD, Seng C, Howe TS, Koh SB, Abdullah HR. Preoperative anemia, functional outcomes, and quality of life after hip fracture surgery. *J Am Geriatr Soc*. 2018;66(8):1524–31.
- Hagino T, Ochiai S, Sato E, Maekawa S, Wako M, Haro H. The relationship between anemia at admission and outcome in patients older than 60 years with hip fracture. *J Orthop Traumatol*. 2009;10(3):119–22.
- Ng TP, Feng L, Nyunt MS, Larbi A, Yap KB. Frailty in older persons: multi-system risk factors and the Frailty Risk Index (FRI). *J Am Med Dir Assoc*. 2014;15(9):635–42.
- Bell KM, Bechara BP, Hartman RA, Shively C, Frazier EC, Lee JY, et al. Influence of number of operated levels and postoperative time on active range of motion following anterior cervical decompression and fusion procedures. *Spine (Phila Pa 1976)*. 2011;36(4):263–8.
- Ohnishi K, Miyamoto K, Kato T, Shimizu K. Effects of wearing halo vest on gait: three-dimensional analysis in healthy subjects. *Spine (Phila Pa 1976)*. 2011;30(7):750–5.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

