

CORRECTION

Open Access



Correction: Comparative intra- and inter-observer reliability of two methods for evaluating intraoperative ultrasonography-based spinal cord hyperechogenicity intensity in degenerative cervical myelopathy

Huachuan Wu^{1,2*†}, Guoliang Chen^{1,2†}, Xianlong Li^{1,2}, Zhengya Zhu¹, Zuofeng Xu³, Xizhe Liu^{2*†} and Shaoyu Liu^{1,2}

Correction: *BMC Musculoskeletal Disord* 23, 630 (2022)
<https://doi.org/10.1186/s12891-022-05517-0>

Following the publication of the original article [1] the authors reported an error in the 2nd paragraph of section “Study population”.

“Their mean age at surgery was 60.8 ± 10.3 years and the average symptom duration was 40.7 ± 34.1 months.”

Should be corrected to:

“Their mean age at surgery was 61.2 ± 10.8 years and the average symptom duration was 23.36 ± 22.11 months.”

The original article [1] has been updated.

Author details

¹Department of Orthopaedic Surgery, Innovation Platform of Regeneration and Repair of Spinal Cord and Nerve Injury, The Seventh Affiliated Hospital, Sun Yat-Sen University, Shenzhen 518107, China. ²Orthopaedic Research Institute/Department of Spinal Surgery, Guangdong Provincial Key Laboratory of Orthopedics and Traumatology, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510080, China. ³Department of Ultrasound, The Seventh Affiliated Hospital of Sun Yat-Sen University, Shenzhen, China.

Published online: 06 September 2022

Reference

1. Wu H, Chen G, Li X, et al. Comparative intra- and inter-observer reliability of two methods for evaluating intraoperative ultrasonography-based spinal cord hyperechogenicity intensity in degenerative cervical myelopathy. *BMC Musculoskeletal Disord.* 2022;23:630. <https://doi.org/10.1186/s12891-022-05517-0>.

The original article can be found online at <https://doi.org/10.1186/s12891-022-05517-0>.

[†]Huachuan Wu and Guoliang Chen contributed equally to this work and should be considered co-first authors.

[†]Xizhe Liu and Huachuan Wu contributed equally to this work and should be considered co-corresponding author, with Huachuan Wu as the first corresponding author.

*Correspondence: wuhch6@mail2.sysu.edu.cn; gzxzhliu@qq.com

²Orthopaedic Research Institute/Department of Spinal Surgery, Guangdong Provincial Key Laboratory of Orthopedics and Traumatology, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou 510080, China
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.