

CORRECTION

Open Access



Correction: Early detection and intervention for acute perforated peptic ulcer after elective spine surgeries: a review of 13 cases from 24,026 patients

Tung-Yi Lin^{1*}, Yu-Chun Chuang^{1†}, Fu-Cheng Kao², Chiu Ping-Yeh², Tsung-Ting Tsai², Tsai-Sheng Fu¹ and Po-Liang Lai²

Correction: *BMC Musculoskeletal Disorders* 22, 548 (2021)
<https://doi.org/10.1186/s12891-021-04443-x>

Following publication of the original article [1], thirteen patients experienced an acute perforated peptic ulcer after elective spine surgery at our hospital between January 2000 and September 2016.

The original article [1] has been updated.

Reference

1. Lin TY, Chuang YC, Kao FC, et al. Early detection and intervention for acute perforated peptic ulcer after elective spine surgeries: a review of 13 cases from 24,026 patients. *BMC Musculoskeletal Disorders*. 2021;22:548. <https://doi.org/10.1186/s12891-021-04443-x>.

Published online: 19 January 2023

[†]Tung-Yi Lin and Yu-Chun Chuang equal contribution as first author.

The original article can be found online at <https://doi.org/10.1186/s12891-021-04443-x>.

*Correspondence:

Tung-Yi Lin
ross_1222@hotmail.com

¹ Department of Orthopaedic Surgery, Chang Gung Memorial Hospital, Keelung branch, and Chang Gung University College of Medicine, Taoyuan, Taiwan, No. 222 Mai-King Road, Keelung, Taiwan

² Department of Orthopaedic Surgery, Spine Section, Bone and Joint Research Center, Chang Gung Memorial Hospital, Linkou branch, and Chang Gung University College of Medicine, Taoyuan, Taiwan

