ERRATUM

Open Access



Erratum to: Clinical trial and in-vitro study comparing the efficacy of treating bony lesions with allografts versus synthetic or highly-processed xenogeneic bone grafts

Eva Johanna Kubosch¹, Anke Bernstein¹, Laura Wolf¹, Tobias Fretwurst², Katja Nelson² and Hagen Schmal^{1,3,4*}

Erratum

After publication of the original article [1], the authors noticed an error in the 'Methods' section. Within the 'Patients' sub-section, the first item in the list of synthetic or highly processed xenogeneic bone substitutes (SBS) should have been given as Orthoss[®], rather than BioOss[®]. The sentence should therefore have read as follows:

"The following SBS were used: Orthoss" (Geistlich, Wolhusen, Switzerland) in 41.4 %, and at clearly lower percentages, respectively: Norian SRS" (Synthes) (8.6 %), Chronos" (Synthes) (1.7 %), Atlantik" (Argomedical) (2.6 %), Alaska" (Argomedical) (0.9 %), Endobone" (Biomet) (4.3 %), Pyrost" (Stryker) (0.9 %), Nanostim" (Medtronic) (37.1 %), Actifuse" (Baxter) (0.9 %), Tutobone" (Novomedics) (0.9 %) and PerOssal" (Botiss) (0.9 %)."

Author details

¹Department of Orthopedics and Trauma Surgery, Albert-Ludwigs University Medical Center, Freiburg, Germany. ²Department of Craniomaxillofacial Surgery, Albert-Ludwigs University Medical Center, Freiburg, Germany. ³Department of Orthopaedics and Traumatology, Odense University Hospital, Odense, Denmark. ⁴Department of Clinical Research, University of Southern Denmark, Odense, Denmark.

Published online: 09 March 2016

Reference

 Kubosch EJ, Bernstein A, Wolf L, Fretwurst T, Nelson K, Schmal H. Clinical trial and in-vitro study comparing the efficacy of treating bony lesions with allografts versus synthetic or highlyprocessed xenogeneic bone grafts. BMC Musc Disord. 2016;17:77.

* Correspondence: hagen.schmal@freenet.de

³Department of Orthopaedics and Traumatology, Odense University Hospital, Odense, Denmark



© 2016 Kubosch et al. **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

¹Department of Orthopedics and Trauma Surgery, Albert-Ludwigs University Medical Center, Freiburg, Germany