

CORRECTION

Open Access



Correction to: Musculoskeletal disorders and their associations with health- and work-related factors: a cross-sectional comparison between Swedish air force personnel and army soldiers

Matthias Tegern^{1,2*}, Ulrika Aasa¹, Björn O. Äng^{2,3,4} and Helena Larsson^{2,5}

Correction to: BMC Musculoskeletal Disorders 21, 303 (2020)
<https://doi.org/10.1186/s12891-020-03251-z>

Following publication of the original article [1], the authors noticed that the numbers 1–5 in the Methods section under subsection “Health- and work-related questions” were mistakenly interpreted as references.

Health- and work-related questions

“To assess their self-rated general health, the following questions were used: How do you experience your [1] “physical health” [2] “mental health” [3] “social environment” [4] “physical environment” and [5] “work ability”? A seven-point scale with answers ranging from “very poor” to “excellent, cannot be better” was used. In the analyses, the answers were collapsed and coded into: poor (≤ 3), good [4, 5], or excellent (≥ 6) according to Larsson et al. [16].”

The original article [1] has been updated.

Author details

¹Department of Community Medicine and Rehabilitation, Unit of Physiotherapy, Umeå University, Umeå, Sweden. ²Department of Neurobiology, Care Sciences and Society, Division of Physiotherapy, Karolinska Institutet, Huddinge, Sweden. ³School of Education, Health and

Social Studies, Dalarna University, Falun, Sweden. ⁴Centre for Clinical Research Dalarna, Uppsala University, Falun, Sweden. ⁵Swedish Armed Forces, HQ, Stockholm, Sweden.

Published online: 10 October 2020

Reference

1. Tegern M, Aasa U, Äng BO, et al. Musculoskeletal disorders and their associations with health- and work-related factors: a cross-sectional comparison between Swedish air force personnel and army soldiers. *BMC Musculoskeletal Disorders*. 2020;21:303. <https://doi.org/10.1186/s12891-020-03251-z>.

The original article can be found online at <https://doi.org/10.1186/s12891-020-03251-z>.

* Correspondence: matthias.tegern@umu.se

¹Department of Community Medicine and Rehabilitation, Unit of Physiotherapy, Umeå University, Umeå, Sweden

²Department of Neurobiology, Care Sciences and Society, Division of Physiotherapy, Karolinska Institutet, Huddinge, Sweden

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.